



Statistical testing and power analysis for brain-wide association study

Weikang Gong^{a,b}, Lin Wan^{c,b}, Wenlian Lu^{d,e,f}, Liang Ma^{g,b}, Fan Cheng^{f,e}, Wei Cheng^e, Stefan Grünewald^{a,b}, Jianfeng Feng^{e,d,f,h,*}



^aKey Laboratory of Computational Biology, CAS-MPG Partner Institute for Computational Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China

^bUniversity of Chinese Academy of Sciences, Beijing 100049, China

^cNCMIS, LSC, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China

^dSchool of Mathematical Sciences, Fudan University, Shanghai 200433, China

^eInstitute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai 200433, China

^fShanghai Center for Mathematical Sciences, Fudan University, Shanghai 200433, China

^gBeijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China

^hDepartment of Computer Science, University of Warwick, Coventry CV4 7AL, UK

ARTICLE INFO

Article history:

Received 30 May 2017

Revised 7 January 2018

Accepted 27 March 2018

Available online 5 April 2018

Keywords:

Brain-wide association study

Random field theory

Functional connectivity

Statistical power

ABSTRACT

The identification of connexel-wise associations, which involves examining functional connectivities between pairwise voxels across the whole brain, is both statistically and computationally challenging. Although such a connexel-wise methodology has recently been adopted by brain-wide association studies (BWAS) to identify connectivity changes in several mental disorders, such as schizophrenia, autism and depression, the multiple correction and power analysis methods designed specifically for connexel-wise analysis are still lacking. Therefore, we herein report the development of a rigorous statistical framework for connexel-wise significance testing based on the Gaussian random field theory. It includes controlling the family-wise error rate (FWER) of multiple hypothesis testings using topological inference methods, and calculating power and sample size for a connexel-wise study. Our theoretical framework can control the false-positive rate accurately, as validated empirically using two resting-state fMRI datasets. Compared with Bonferroni correction and false discovery rate (FDR), it can reduce false-positive rate and increase statistical power by appropriately utilizing the spatial information of fMRI data. Importantly, our method bypasses the need of non-parametric permutation to correct for multiple comparison, thus, it can efficiently tackle large datasets with high resolution fMRI images. The utility of our method is shown in a case-control study. Our approach can identify altered functional connectivities in a major depression disorder dataset, whereas existing methods fail. A software package is available at <https://github.com/weikanggong/BWAS>.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

The human brain connectome is usually modeled as a network. In the brain's network, accurately locating the connectivity variations associated with phenotypes, such as clinical symptoms, is critical for neuroscientists. With the development of neuroimaging technology and an increasing number of publicly available datasets, such as the 1000 Functional Connectomes Project (FCP) (Biswal et al., 2010), Human Connectome Project (HCP) (Glasser et al., 2016) and UK-Biobank (Miller et al., 2016), large-

scale, image-based association studies have become possible and should help us improve our understanding of human brain functions.

Using a priori knowledge of brain parcellation (e.g. Automated Anatomical Labeling atlas, Rolls et al., 2015) or an adoption of data-driven parcellation (e.g. Independent Component Analysis, Beckmann and Smith, 2004) to analyze the human connectome is the most popular approach, and many statistical methods have been designed for them (Zalesky et al., 2012; Kim et al., 2014). However, methods that designed specifically for voxel-level connectivity analysis are still lacking. Therefore, in this paper, a statistical framework for brain-wide association study (BWAS) is proposed (Cheng et al., 2015a; 2015b; 2016). It directly uses voxels as nodes to define brain networks, and then tests the associations of each functional connectivity with phenotypes.

* Corresponding author at: Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai 200433, China.

E-mail address: jianfeng64@gmail.com (J. Feng).

To conduct a systematic, fully-powered BWAS, two main issues should be carefully addressed. First, a multiple correction method to control the false-positive rate of massive univariate statistical tests should be developed. Second, a power analysis method to estimate the required sample size should be designed. One may ask whether the methods widely used in region-level studies can be directly generalized to connexel-level studies. Two issues hinder such a direct generalization. First, the statistical tests have more complex spatial structures in BWAS. Therefore, as shown in our analysis, some widely-used multiple correction methods, which do not utilize the spatial information of data (e.g. Bonferroni correction and false discovery rate (FDR), Benjamini and Hochberg, 1995; Benjamini and Yekutieli, 2001), may not be powerful enough to detect signals. Second, although non-parametric permutation methods (Nichols and Holmes, 2002) may account for the complex structures among hypothesis tests to provide a valid threshold, they are computationally very expensive in connexel-wise studies, owing to the requirement of performing billions of statistical tests. Therefore, an accurate and efficient method for multiple comparison problem and power analysis is needed.

Random field theory (RFT) is an important statistical tool in brain image analysis, and it has been widely used in the analysis of task fMRI data (Penny et al., 2011) and structure data (Ashburner and Friston, 2000). Statistical parametric maps (SPM) are usually modeled as a discrete sampling of smooth Gaussian or related random fields (Penny et al., 2011). The random field theory can control the FWER of multiple hypothesis testings by evaluating whether the observed test statistic, or the spatial extent of clusters exceeding a cluster-defining threshold (CDT), is large by chance, which is known as peak-level and cluster-level inference respectively. Since Adler's early work on the geometry of random field (Adler, 1981; Adler and Taylor, 2009), theoretical results for different types of random fields have been obtained, such as Gaussian random field (Friston et al., 1994; Worsley et al., 1996), t , χ^2 , F random fields (Worsley, 1994; Cao, 1999), multivariate random field (Taylor and Worsley, 2008), cross-correlation random field (Cao et al., 1999). Among them, only the cross-correlation field is designed for connectivity analysis. In that framework, the voxel-level functional network is modeled as a six-dimensional cross-correlation random field, and the maximum distribution of the random field is used to identify strong between-voxel connections. Different from the above works, the aim of BWAS is to identify connectivities that are associated with phenotypes. To the best of our knowledge, no previous works have addressed this problem. In this paper, we show that the statistical map of BWAS, under the null hypothesis, can be modeled as a Gaussian random field with a suitable smoothness adjustment. Therefore, topological inference methods, such as peak intensity and cluster extent, are generalized from voxel-wise analysis to functional connectivity analysis.

Besides controlling the type I error rate, estimating power and the required sample size for BWAS are also important. In genetics, for example, a high-quality GWAS analyzing one million single nucleotide polymorphism (SNP) usually requires tens of thousands of samples to reach adequate statistical power. In contrast, previous BWAS analyses of schizophrenia, autism and depression have only had sample sizes less than one thousand (Cheng et al., 2015a; 2015b; 2016). Therefore, compared to GWAS, it is natural to ask if BWAS, which is usually based on a limited sample size, can withstand the rigors of a large number of hypothesis tests. In this regard, most existing power analysis methods are designed for voxel-wise fMRI studies, including, for example, the simulation based method (Desmond and Glover, 2002), the non-central distribution based method (Mumford and Nichols, 2008), and the method based on non-central random field theory (ncRFT) (Hayasaka et al., 2007). Among them, the ncRFT-based method can both take into account the spatial structure of fMRI data and avoid time con-

suming simulation. Therefore, to analyze the power of BWAS, we adopted a methodology similar to that of the ncRFT-based method (Hayasaka et al., 2007). The signals at functional connectivities are modeled as a non-central Gaussian random field, and the power is estimated by a modified Gaussian random field theory.

In this paper, a powerful method to address the multiple comparison problem is proposed for BWAS (Fig. 1). This method uses Gaussian random field theory to model the spatial structure of voxel-level connectome. It can test the odds that either the effect size of every single functional connectivity (peak-level inference) or the spatial extent of functional connectivity clusters exceeding a cluster-defining threshold (cluster-level inference) is large by chance. The performance of the method is tested in two resting-state fMRI datasets, and in both volume-based and surface-based fMRI data. Our method can control the false-positive rate accurately. Compared with Bonferroni correction and false discovery rate (FDR) approaches, our method can achieve a higher power and filter out false-positive connectivities by utilizing the spatial information. In addition, we develop a method to approximate the power of peak-level inference by a modified Gaussian random field theory (Fig. 2). Power can be estimated in any specific location of connectome efficiently, which can help to determine the sample size for BWAS. The utility of our method is shown by identifying altered functional connectivities and estimating the required sample sizes in major depression disorder. The software package for BWAS can be downloaded at <https://github.com/weikanggong/BWAS>.

2. Material and methods

2.1. Connexel-wise general linear model

The popular general linear model approach is used in BWAS. Briefly, a voxel-level functional network is estimated for each subject using the fMRI data, and the association between each functional connectivity and phenotype of interest is tested using the general linear model.

In detail, the individual functional network is constructed first by calculating the Pearson correlation coefficients (PCC) between every pair of voxel time series. Let m be the number of voxels, s be the subject, and $R^{(s)} = [r_{ij}^{(s)}]_{m \times m}$ be the $m \times m$ functional network matrix for subject s . Each element of $R^{(s)}$ is the correlation coefficient between voxel time series i and j for subject s . An element-wise Fisher's Z transformation is then applied as $Z^{(s)} = [z_{ij}^{(s)}]_{m \times m} = \left[\frac{1}{2} \log \left(\frac{1+r_{ij}^{(s)}}{1-r_{ij}^{(s)}} \right) \right]_{m \times m}$, so that $z_{ij}^{(s)}$ will approximate a normal distribution. For every functional connectivity, a general linear model (GLM) is fitted by

$$Y_{ij} = XB_{ij} + \epsilon_{ij}$$

where, $Y_{ij} = (z_{ij}^{(1)}, z_{ij}^{(2)}, \dots, z_{ij}^{(n)})$ is an $n \times 1$ vector of functional connectivities between voxel i and j across n subjects, X is the common $n \times q$ design matrix, $B_{ij} = (\beta_{ij}^1, \beta_{ij}^2, \dots, \beta_{ij}^q)$ is a $q \times 1$ vector of regression coefficients, and ϵ_{ij} is an $n \times 1$ vector of random error, which is assumed to be an independent and identically distributed Gaussian random variable $N(0, \sigma_{ij}^2)$ across subjects. The ordinary least square estimator for B_{ij} is $\hat{B}_{ij} = (X'X)^{-1}X'Y_{ij}$, and for σ_{ij}^2 , it is $\hat{\sigma}_{ij}^2 = (Y_{ij} - X\hat{B}_{ij})'(Y_{ij} - X\hat{B}_{ij})/(n - q)$. Then, a Student's t -statistic at functional connectivity between voxel i and j can be expressed as:

$$T_{ij} = \frac{\mathbf{c}\hat{B}_{ij}}{(\mathbf{c}(X'X)^{-1}\mathbf{c}'\hat{\sigma}_{ij}^2)^{\frac{1}{2}}}$$




Fig. 1. A flow chart of brain-wide association study. First, we estimate the voxel-level brain network for each individual. Then, we perform connectivity-wise statistical tests to test the association between each functional connectivity and a phenotype of interest. Finally, peak- and cluster-inference approaches are used to identify significant signals. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)




Fig. 2. Power analysis for brain-wide association study. To estimate power, we first calculate the FWER-corrected threshold Z_α of peak-level inference, and then estimate the effect size γ from a prior statistical map of BWAS. For a target sample size n , and a smoothness level FWHM, we can estimate the power using the random field theory. The power is defined as the probability of finding at least one true-positive signal in a region, in which the false-positive rate α is controlled at a certain level in the whole search region. Finally, the power under different sample sizes and smoothness levels can be estimated iteratively. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

where \mathbf{c} is a $1 \times q$ contrast vector. In BWAS, let β_{ij}^1 be the primary variable of interest, and $\beta_{ij}^2, \dots, \beta_{ij}^q$ be the nuisance covariates included in the regression model. The contrast $\mathbf{c} = (1, 0, \dots, 0)$ will be used to test the hypothesis $\beta_{ij}^1 = 0$, and the T_{ij} -statistics will reflect the significance of the primary variable. Other contrasts can also be used depending on the study design. Finally, the Student's t random variable at each functional connectivity T_{ij} is transformed

to a Gaussian random variable Z_{ij} by transforming t -statistics to p -values and then to Z -statistics.

After the above steps, the connexel-wise Z -statistics form a statistical parametric map in a six-dimensional Euclidian space. The reason is that the spatial location of each Z statistic (or functional connectivity) can be uniquely represented by the coordinates of its two endpoints, each of which is a voxel in a three-dimensional space. Therefore, the structure of the statistical map can be mod-

eled by the random field theory, and the topological inference methods for multiple hypothesis testings are developed in the subsequent section.

2.2. Multiple comparison correction using topological inference methods

2.2.1. Peak-level inference

Peak-level inference controls FWER among multiple hypothesis testings on functional connectivities, i.e., the probability of finding at least one false-positive functional connectivity is controlled under certain level α . It is assumed, under the null hypothesis, that the statistical parametric map of BWAS is a discrete sampling of smooth and stationary Gaussian random field with mean zero and variance one. To control the FWER of multiple hypothesis testings, the maximum distribution of the random field should be known. In our paper, its tail distribution is approximated by the expected Euler characteristic (EC) of the excursion set of random field (Adler and Taylor, 2009; Worsley et al., 1996). The detailed derivation is given in the appendix section. We sketch an overview of the result here.

Let $Z(p, q)$, $p \in \mathcal{P}$, $q \in \mathcal{Q}$ be a $(P+Q)$ -dimensional Gaussian random field spanned by a P -dimensional random field \mathcal{P} and a Q -dimensional random field \mathcal{Q} . At high threshold z_0 , its maximum distribution has a general form:

$$\begin{aligned} \alpha &= P(\max Z(p, q) > z_0) \approx \mathbb{E}(EC) \\ &= \sum_{d=0}^{P+Q} \mu_d(\mathcal{P} \times \mathcal{Q}) \rho_d^Z(z_0) \\ &= \sum_{i=0}^P \sum_{j=0}^Q \mu_i(\mathcal{P}) \mu_j(\mathcal{Q}) \rho_{i+j}^Z(z_0) \end{aligned} \quad (1)$$

where the $\mu_d(\cdot)$ is the d -th dimensional intrinsic volume of the random field, and $\rho_d^Z(z_0)$ is the d -th dimensional EC-density for the Gaussian random field at threshold z_0 ($z_0 > 0$). The methods for calculating $\mu_d(\cdot)$ and $\rho_d^Z(\cdot)$ are shown in Appendix B and a proof of Eq. (1) is shown in Appendix D. Therefore, the α -level FWER-corrected threshold z_0 can be found using Eq. (1), and for one-tailed tests, functional connectivities with Z -values larger than z_0 (or smaller than $-z_0$) are declared as significant.

For different kinds of BWAS analysis, P and Q in Eq. (1) can take different values. For example, for the widely-used volume-based fMRI data, we use $P=Q=3$. If the connectivities are estimated between pairwise vertices on cortical surface, we use $P=Q=2$, and if the connectivities are estimated between subcortical structures and cortical surface, we use $P=3$ and $Q=2$. The estimated FWER-corrected threshold is usually less conservative than Bonferroni correction method, because the intrinsic volume $\mu_d(\cdot)$ in Eq. (1) takes into account both the number of hypothesis tests performed and the correlations among tests, and an increasing of spatial smoothness can make the FWER-corrected threshold z_0 lower. For BWAS, Eq. (1) can be approximately estimated using the results of Gaussian random field, provided that the spatial smoothness is estimated correctly. The reason is that the statistical map of BWAS is generated by a series of non-linear transformation of original fMRI images. As a result, we calculate Eq. (1) as:

$$\begin{aligned} \alpha &= P(\max Z(p, q) > z_0) \\ &\approx \sum_{i=0}^P \sum_{j=0}^Q \mu_i(\mathcal{P}) \mu_j(\mathcal{Q}) \frac{(2\pi)^{-\frac{i+j+1}{2}} (4 \log 2)^{\frac{i+j}{2}}}{\text{FWHM}_z^{i+j}} \\ &\times e^{-\frac{z_0^2}{2}} \sum_{k=0}^{\lfloor \frac{i+j-1}{2} \rfloor} (-1)^k \frac{(2k)!}{k! 2^k} \binom{i+j-1}{2k} z_0^{d-1-2k} \end{aligned} \quad (2)$$

where FWHM_z is the adjusted full-width at half maximum of the Gaussian smooth kernel, which is a function of the original smoothness of fMRI images whose estimation approach is shown in Appendix C. A proof of the Eq. (2) is shown in Appendix E.

2.2.2. Cluster-level inference

Cluster-level inference is also popular in brain image analysis. Here, inference is based on the observed cluster size exceeding certain cluster-defining threshold (CDT) (Friston et al., 1994). We are usually interested in whether the observed cluster size is large by chance, i.e., where the size is on the upper tail of the distribution of maximum cluster size under the null hypothesis. We show that, similar to the voxel clusters in three-dimensional space, the functional connectivity cluster (FC cluster) can also be defined rigorously. Its size can be used as a test statistic for statistical inference, and it has a clear interpretation.

A voxel cluster is a set of spatially connected voxels. To define a FC cluster, we first illustrate the neighborhood relationship between two functional connectivities. Let the endpoints of two functional connectivity be (x_1, y_1) and (x_2, y_2) , if their endpoints are non-overlapped voxels, then two functional connectivities are neighbors if both x_1, x_2 and y_1, y_2 are spatially adjacent voxels. If they share a same endpoint (e.g. $x_1 = x_2$), then they are neighbors if y_1, y_2 are spatially adjacent voxels. Some examples of FC neighbors are shown in Fig. 3(A). Now, consider an undirected graph \mathcal{G} with k nodes, where the nodes are k functional connectivities and two nodes are connected if they are neighbors, then these k functional connectivities form a FC cluster if they form a connected component in the graph \mathcal{G} . Some examples of FC clusters are shown in Fig. 3(B). There are five voxel clusters A, B, C, D, E in a two-dimensional image. The FCs between AB, BC and AD are different FC clusters, and FCs within voxel cluster E also form a single FC cluster. An algorithm for finding FC clusters can be implemented based on the above definition. In our analysis, We use Dulmage–Mendelsohn decomposition to find connected components in graph \mathcal{G} .

Based on the normality and stationarity assumption as in peak-level inference, we propose to use Gaussian random field theory to approximate the null distribution of maximum cluster size. In brief, let M be the number of FCs with z -values exceeding a pre-specified CDT z_0 , N be the number of FC clusters, and S be size of a FC cluster. Suppose that separate FC clusters are independent, then the distribution of maximum cluster size S_{max} for Gaussian random field is (Adler, 1981; Friston et al., 1994):

$$P(S_{max} > s) = 1 - \exp[-\mathbb{E}(N)P(S > s)]$$

The expected number of FC clusters $\mathbb{E}(N)$ at a high CDT z_0 can be approximated by the expected EC of Gaussian random field using Eq. (2):

$$\mathbb{E}(N) \approx \mathbb{E}(EC) = \sum_{i=0}^P \sum_{j=0}^Q \mu_i(\mathcal{P}) \mu_j(\mathcal{Q}) \rho_{i+j}^Z(z_0)$$

The distribution of S can be approximated by (Adler, 1981; Nosko, 1969):

$$P(S > s) = \exp \left[- \left(\frac{\Gamma((P+Q)/2 + 1) \mathbb{E}(N) s}{\mathbb{E}(M)} \right)^{\frac{2}{(P+Q)}} \right]$$

where $\mathbb{E}(M) = m[1 - \Phi(z_0)]$, and m is the number of functional connectivities, and $\Phi(\cdot)$ is the cumulative distribution function of standard normal distribution. The above theory is a generalization of previous result (e.g. Friston et al., 1994 and Hayasaka and Nichols, 2003), except the use of Eq. (2) to approximate the expected number of clusters $\mathbb{E}(N)$.

Therefore, in the cluster-level inference, small-sized FC clusters are more likely to be identified as false positives and filtered out




Fig. 3. Two-dimensional diagrams of FC neighbors and FC clusters. The size of FC clusters exceeding a cluster-defining threshold is used as a test statistic in the cluster-level inference. (A) In BWAS, there are two cases that FCs are neighbors. In situation 1, two FCs share a common endpoint and another two endpoints are spatial neighbors. In situation 2, two pairs of endpoints of two FCs are all spatial neighbors. Situation 3 is a special case of situation 2. (B) In BWAS, FCs can form a cluster in two ways. The first one is the FCs between voxel cluster AB, BC and AD, and the second one is FCs within a voxel cluster E. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

(e.g. the red link in Fig. 3(B)). Large-sized FC clusters represent an existence of association signals either between two different voxel clusters (e.g. the blue and green ones in Fig. 3(B)) or within a single voxel cluster (e.g. the yellow one in Fig. 3(B)). For example, consider that one performed a case-control BWAS, then the identified FCs can be either altered connections between two brain regions or within a single region.

2.3. Validating peak- and cluster-level inference in real data

2.3.1. Data

We briefly illustrate the data used in this article, all details are provided in Appendix A.

Two resting-state fMRI datasets are used in our analysis: (1) 197 subjects from the Cambridge dataset in the 1000 Functional Connectomes Project (1000 FCP); (2) 222 subjects from the Southwest University (SWU) dataset in the International Data-sharing Initiative (IDNI). The subjects in the two datasets are all healthy people with similar demographic information. They are preprocessed using standard preprocessing pipelines implemented in Data Processing and Analysis for Brain Imaging (DPABI) (Yan et al., 2016). Finally, All fMRI data are registered to $3 \times 3 \times 3$ mm³ standard space, and 47,636 voxel time series within each subject's 90 cerebrum regions (based on AAL template, Rolls et al., 2015) are extracted. They are then smoothed by 3D Gaussian kernels with FWHM = 0, 2, 4, 6, 8, 10, 12 mm on each dimension.

In addition, the above data are also mapped on to the Conte69 surface-based atlas using the Connectome Workbench software. They are smoothed by 2D Gaussian kernels restricted on the cortical surface with FWHM = 0, 4, 8 mm. Finally, 32,492 vertex time series on the left cortical surface are used in our analysis.

2.3.2. Evaluating the proposed methods

To evaluate whether the random field theory can actually control the FWER in real data analysis (including connexel-wise FWER and cluster-size FWER), we compared our method with empirical permutation results in real data. Similar approaches have previously been adopted to validate the random field theory in task-activation studies (Eklund et al., 2016; 2012).

The following procedures were carried out in each of the volume-based and surface-based fMRI datasets. First, subjects were randomly divided into two groups. Second, BWAS was performed to compare the whole-brain functional connectivities between two groups (approximately 1.13×10^9 connections in volume-based data, and 5.28×10^8 connections in surface-based data). The peak-

and cluster-level inference approaches were applied to find significant signals. Third, the above two steps were repeated 2000 times. FWER was then estimated by computing that proportion of permutations in which any significant signal was found. The signal was functional connectivities in the peak-level inference, and FC clusters exceeding certain CDT in the cluster-level inference. Since subjects were all healthy people with similar demographic information, and their group labels were randomly assigned, we expected that there were no group differences. Therefore, if the proposed approach is valid, the proportion of analysis with at least one significant effect should be close to the nominal error rate 0.05.

2.4. Comparing with other multiple correction methods

We compared our proposed method with connexel-wise Bonferroni correction and false discovery rate (FDR-BH, Benjamini and Hochberg, 1995; FDR-BY, Benjamini and Yekutieli, 2001) in terms of the observed power and false-discovery rate. To mimic real data, we did not use completely simulated data, but rather, we adopted a widely used evaluation methodology in GWAS (e.g. Yang et al., 2014; Zhou and Stephens, 2012), which directly add simulated signals to real data. The data used here were 197 subjects in the Cambridge dataset in four smoothness levels (FWHM = 0,4,8,12 mm).

2.4.1. Simulation procedures

First, two cerebrum regions within the AAL template were randomly selected. BOLD signals of voxels within these two regions were extracted. Second, subjects were randomly divided into two groups. For each subject in one group, a common signal was added to a small cluster of voxel time series of each region. Therefore, the functional connectivities between voxels in these two clusters were increased, which formed a single FC-cluster. Third, a two-sample *t*-test was used to compare two groups of functional connectivities. Five methods, including Bonferroni correction, FDR-BH (Benjamini and Hochberg, 1995), FDR-BY (Benjamini and Yekutieli, 2001), peak-level inference, and cluster-level inference (with different CDT), were used to control the false-positive rate of multiple hypothesis testings. The steps two and three were repeated for 100 times.

Four free parameters were found in our simulation: (1) voxels selected from real data, (2) signal width, i.e., the number of altered functional connectivities, (3) effect size of the signal and (4) image smoothness. In Section 3, we will report the results of comparisons among the different combinations of parameters.

2.4.2. Performance metrics

Two metrics were used to evaluate the performance: the observed power and false-discovery rate. The observed power was calculated as the number of discovered true-positive functional connectivities divided by the total number of true-positive connectivities. The observed false-discovery rate was calculated as the number of discovered false-positive functional connectivities divided by total number of discovered functional connectivities. In Section 3, we will report the average power and false-discovery rate of different methods.

2.5. Statistical power analysis

A method to estimate the statistical power of peak-level inference is proposed (Fig. 2). Power is defined as the probability of finding at least one true-positive signal for a region (denoted as B) in which the false-positive rate α is controlled at a certain level in the whole search region (denoted as A) (Friston et al., 1994). To estimate power, four parameters should be specified: (1) the threshold that controls connectivity-wise FWER α , (2) the effect size of true signal γ , (3) the sample size n , and (4) the smoothness of image.

First, if we assume that the primary variable of interest, β_{ij}^1 , is subject to a normal distribution $N(\mu_{ij}, \sigma_{ij}^2)$, the null hypothesis is $H_0 : \mathbb{E}(\beta_{ij}^1) = 0$. Therefore, every test statistic Z_{ij} is subject to $N(0, 1)$. The whole search region A is a central Gaussian random field with mean zero and variance one at each point. The threshold z_0 to control the FWER at α is obtained by the random field theory (Eq. (2)):

$$\alpha = P(\max_{(p,q) \in A} Z(p, q) > z_0 | H_0)$$

where (p, q) are the coordinates of the functional connectivities.

Then, under the alternative hypothesis $H_1 : \mathbb{E}(\beta_{ij}^1) = \mu_{ij}$, the test statistics Z_{ij} is subject to $N(\sqrt{n}\mu_{ij}/\sigma_{ij}, 1)$, where n is the sample size. The $\gamma_{ij} = \mu_{ij}/\sigma_{ij}$ is called effect size at FC_{ij} . We further assume that the distribution of signals will be the same in region B, i.e., all β_{ij}^1 is subject to the same normal distribution $N(\mu, \sigma^2)$. Therefore, region B is a non-central Gaussian random field $Z^*(p, q)$ with mean $\sqrt{n}\gamma$ and variance one at each point. The power in the search region $B \subset A$ can be expressed as:

$$\text{Power} = P(\max_{(p,q) \in B} Z^*(p, q) > z_0 | H_1)$$

The non-central Gaussian random field $Z^*(p, q)$ can be transformed to a central Gaussian random field by the following element-wise transformation:

$$Z(p, q) = Z^*(p, q) - \sqrt{n}\gamma$$

Therefore, the power in region B can still be calculated using Eq. (2):

$$\text{Power} = P(\max_{(p,q) \in B} Z(p, q) > z_0 - \sqrt{n}\gamma | H_0)$$

Three issues remain. The first involves selecting region B. When estimating power, we select region B as consisting of functional connectivities between two three-dimensional balls, with the diameter of each ball being equal to the intrinsic FWHM of the image (Fig. 2). The idea is that signals within such ball are usually homogeneous as a result of smoothing the images.

The second issue involves the random field theory which can only approximate the right tail of the maximum distribution. Therefore, the above method may lead to an inaccurate estimation when $z_0 - \sqrt{n}\gamma$ is small. To address this problem, we propose to use the following heuristic modification:

$$\text{Power} = 1 - \exp \left[-P(\max_{(p,q) \in B} Z(p, q) > z_0 - \sqrt{n}\gamma | H_0) \right]$$

This formula ensures that the power is between zero and one, which shows excellent performance in the simulation.

The last issue concerns estimating the effect size, which is typically estimated from the statistical map of a pilot study using the same study design. Suppose that the pilot BWAS study used n^* samples. Then, the estimated effect size at FC_{ij} is (Joyce and Hayasaka, 2012):

$$\hat{\gamma}_{ij} = Z_{ij} / \sqrt{n^*}$$

Using the above formula, the power can be estimated for each functional connectivity to form a power map on six-dimensional space, but it is quite difficult to visualize such a maps. Therefore, to report the power of a study, we estimate the effect size of every FC to form an empirical distribution. The power curves of different sample sizes and effect sizes under certain power (e.g. 90% power) are analyzed and reported.

2.5.1. Simulation-based validation for power analysis

To test whether the proposed method can estimate power accurately, we performed a simulation study. Briefly, we simulated a case-control study with known effect size in a subset of functional connectivities, and we compared the observed power and the theoretical power.

In detail, first, we generated two sets of 10,000 three-dimensional independent Gaussian white noise images, with 30 voxels per dimension. Second, the images were smoothed by Gaussian kernels with FWHM ranging from 3 to 6 voxels. Third, a ball was extracted for subsequent analysis with radius being equal to the FWHM of images and located at the center of each image. This guaranteed the uniform smoothness. Fourth, every 20 images were combined to form 500 simulated four-dimensional fMRI data. We denoted the images in the first set as $(A_1, A_2, \dots, A_{500})$ and the images in the second set as $(B_1, B_2, \dots, B_{500})$. Fifth, the Pearson correlation coefficients were calculated between time series of pairwise voxels of images A_i and B_i , and a Fisher's Z transformation was then performed. Sixth, two groups of images from two sets were randomly selected, with each group consisting of 200 samples. A Z-map was then generated by fitting each functional connectivity to a general linear model to compare the two groups. Seventh, signals were then added to functional connectivities between the two balls. Specifically, a signal was the mean intensity difference between two groups. We then estimated power using simulated data under different parameters, including image smoothness FWHM, sample size n and effect size γ (Fig. 2). The steps six and seven were repeated for 10,000 times under each parameter setting, and the maximum statistics were recorded at each simulation. The empirical power was estimated by the proportion of maximum statistics exceeding the FWER-corrected 0.05 threshold. We compared the results of simulation with the proposed theoretical method.

3. Results

3.1. Overview of the proposed approaches

Figs. 1 and 2 show the diagrams of the proposed approaches. Fig. 4 shows the multiple comparison threshold of different approaches in a typical BWAS study. In the study, the fMRI data have a spatial resolution of $3 \times 3 \times 3 \text{ mm}^3$. A total of 47,636 voxels in the cerebrum regions were used.

Methods that control connectivity-wise FWER, including Bonferroni correction and peak-level inference, provide evidence of association of each individual functional connectivities that survive the threshold. Bonferroni correction is always the most conservative one. The peak-level inference is more powerful when the smoothness of images increased. As shown in Fig. 4, the FWER-corrected

Table 1

The estimated smoothness of different datasets used in our analysis (FWHM in mm).

Applied smoothness		0 mm	2 mm	4 mm	6 mm	8 mm	10 mm	12 mm
Estimated smoothness	Cambridge volume-fMRI	4.5 mm	4.6 mm	7 mm	10.6 mm	13.2 mm	15.9 mm	17.6 mm
	Cambridge surface-fMRI	4.7 mm		6.9 mm		11.1 mm		
	SWU volume-fMRI	5.1 mm	5.2 mm	7.5 mm	10.9 mm	13.4 mm	15.3 mm	17.9 mm
	SWU surface-fMRI	5.2 mm		7.7 mm		12.2 mm		




Fig. 4. A comparison of the multiple comparison thresholds provided by different methods in BWAS using $3 \times 3 \times 3$ mm³ volume-based fMRI data. Methods that control FWER (Bonferroni correction and peak-level inference) provide universal thresholds across FCs. The threshold of FDR approaches depend on the rank of p -values of FCs. The cluster-defining threshold of cluster-level inference is an universal threshold across FCs, and a subsequent correction on the size of FC clusters is applied. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

threshold can be 1 to 2 order of magnitudes less conservative than Bonferroni correction.

Methods that control connectivity-wise FDR, including FDR-BH and FDR-BY approaches, control the proportion of false-positive findings smaller than a pre-specified level q (e.g. 5%). For the widely-used FDR-BH approach, it compares the i -th smallest p -value $p_{(i)}$ with $\frac{i}{n}q$, where n is the total number of hypothesis tests, and declares the first k smallest p -values as significant that if $p_{(k)}$ is the largest p -value satisfies: $p_{(k)} < \frac{k}{n}q$ (Fig. 4). Therefore, the power of FDR approaches highly depends on the observed p -values, which can be more or less powerful than the peak- and cluster-level inference. For example, controlling the FDR at $q = 0.05$ is very stringent, owing to the billions of statistical tests performed in BWAS. However, it can be more powerful when many of the p -values meet the requirement of the data-driven threshold. A method that controls connectivity-wise FWER can also control connectivity-wise FDR.

The cluster-level inference approach tests the size of the FC clusters exceeding a CDT. A significant FC cluster can provide evidence that there exist association signals somewhere in this FC cluster. None of the individual functional connectivities in the cluster can be declared as significant ones. This approach is usually sensitive to spatially extended signals. Moreover, when the CDT equals the FDR threshold, the connectivity-wise FDR can be controlled, and when the CDT equals the FWER threshold, it is equivalent to control the connectivity-wise FWER.

3.2. Validating peak- and cluster-level inference in real data

3.2.1. Estimated connexel-wise and cluster-size FWER in real datasets

We evaluate whether the proposed method can control the FWER in real data analysis, including the connexel-wise FWER for the peak-level inference and cluster-size FWER for the cluster-level inference. We compared the theoretical FWER with the empiri-

cal FWER estimated by permutation approaches using real data. The experimental procedures are illustrated in Section 2.3.2. For volume-based fMRI data, we used 2 datasets under 7 smoothness levels. For surface-based fMRI data, we used 2 datasets under 3 smoothness levels. The estimated smoothness of different datasets are shown in Table 1.

Fig. 5 shows the estimated FWER of peak- and cluster-level inference methods using volume-based fMRI data. We found that the peak-level approach is valid, as most of the estimated FWERs lie in the binomial confidence interval of 2000 permutations (dashed line). The cluster-level inference is also valid if the CDT is larger than 5. However, when the CDT becomes smaller, the false-positive rate will exceed the nominal level, because the assumptions of the theory may break down.

Fig. 6 shows, for cluster-level inference, the comparison of the estimated cluster-size threshold of random field theory and permutation approach at low smoothness levels. Different from the above analysis, we directly compare two thresholds because the 95% quantiles of empirical maximum cluster-size distribution cannot be estimated accurately in permutation tests. The reason is that when the smoothness is low, the size of FC clusters is usually small, thus, there are many ties in the maximum cluster-size distribution. A good agreement between the two thresholds demonstrates the validity of cluster-size inference at low smoothness level, and the CDT can even be lower comparing with the above analysis ($Z=4.5$).

Fig. 7 shows the estimated FWER of peak- and cluster-level inference methods using surface-based fMRI data. We found that, when no spatial smoothing is applied (FWHM = 0 mm), our approach is more conservative than permutation approach. The method works well when we smooth the data. However, to the best of our knowledge, there are no standard preprocessing pipelines for surface-based resting-state fMRI data, thus, our surface-mapping approach may not be optimal for BWAS and different preprocessing pipelines may affect the performance of our approach. Therefore, the robustness of the approach should be tested in the future.

3.2.2. Distribution of functional connectivity data

We tested whether functional connectivities data, i.e., Fisher's Z transformed correlation coefficients, were subject to normal distributions, which was a critical assumption for Gaussian random field theory. We performed one-sample Kolmogorov–Smirnov test to test the normality of each functional connectivity in both Cambridge and Southwest University datasets. Supplement Figs. F.13 and F.15 show the results. As most of the p -values are larger than 0.05, we conclude that the normality assumption is met.

3.2.3. The choice of cluster-defining threshold

For cluster-level inference, the expected Euler Characteristics is used to approximate the expected number of clusters in the random field theory, assuming that the absence of holes when CDT is applied. However, this assumption may not be true when the CDT is not high enough or the data are not smooth enough. Therefore, we compared the expected Euler characteristics calculated based on the Gaussian random field theory with the observed expected number of clusters across different levels of CDT in volume-based fMRI data in both Cambridge and Southwest University datasets.




Fig. 5. Validating peak- and cluster-level inference by comparing the theoretical FWER with permutation-based empirical FWER at 0.05. The methods are tested in 2 datasets (top: Cambridge; bottom: SWU) under 7 different smoothness levels (0–12 mm smoothing). The estimated FWER is that proportion of permutations in which any significant signals (functional connectivities or FC clusters) are found by the random field theory. Left: results for peak-level inference. Right: results for cluster-level inference with different clustering-defining thresholds (from 4.9 to 5.9). Almost all the results lie in the binomial 95% confidence interval (the dashed line). (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

The observed expected number of clusters was computed based on an average of 2000 permutations of each dataset. The results are shown in Supplement Figs. F.14 and F.16. We find that, when the applied smoothness is larger than 4 mm, the choice of CDT greater than 5 is very safe for $3 \times 3 \times 3$ mm³ resolution fMRI data to meet the assumption of the random field theory. This is in agreement with our results in the previous section (Fig. 5). When the smoothness is low, we found that there exist a deviation between theory and real data when the CDT is smaller than 5.5. However, the results shown in Fig. 6 indicate the proposed method can provide a valid threshold when the CDT is as low as 4.5 in two datasets.

3.3. Comparing peak- and cluster-level inference with other multiple correction methods

Fig. 8 shows the results of comparisons using 197 subjects in the Cambridge dataset. The experimental procedures are illustrated in Section 2.4. In this analysis, we extracted time series of 306 voxels from the left putamen region and 302 voxels from the left inferior frontal gyrus in each of the 197 subjects, and $306 \times 302 = 92,412$ functional connectivities between these two regions were calculated. Signals were added to a subset of voxel time series, with effect size ranging from 0.5 to 1.5, and smoothness of 0,4,8,12 mm was applied. For cluster-level inference, we used CDT = 3, 3.5 and 4 (Z-value). The following observations are obtained from this simulation:

- Almost all the methods can control false discovery rate in this simulation (below 5%).
- The power of peak-level inference is similar to Bonferroni correction when the smoothness is low (e.g. no spatial smoothing), but it becomes close to FDR-BY and much higher than that

of Bonferroni correction when the smoothness is high (e.g. applied smoothness of 12 mm).

- The false discovery rate and power of cluster-level inference depends on the choice of CDT. The lower the CDT, the higher the false discovery rate and power.
- We can find a CDT whose power is higher than that of the FDR method. In the meantime, the false discovery rate is lower. For example, for cluster-level inference with CDT=3, the power is higher than that of the FDR-BH method, and the false discovery rate is lower. For cluster-level inference with CDT=3.5, the power is higher than that of the FDR-BY method, and the false discovery rate is lower.

Similar results can be obtained by changing the selected voxels and the width of the signal added, as shown in the Supplement Figs. F.17 and F.18. In conclusion, cluster-level inference can increase sensitivity and decrease false-positive rate by filtering out small FC-clusters generated by random noises. Peak-level inference shows increased power when the smoothness is large; thus, it is recommended when performing group-level studies with large applied smoothness.

3.4. Real data analysis: identifying altered functional connectivities in major depression disorder

We applied our method to identify functional connectivity difference between patients with major depression disorder (MDD) and healthy controls. The data used here are part of the data in our previous study (Cheng et al., 2016) which contained 282 patients and 254 demographic information matched controls from Southwest University dataset. We applied BWAS approach to test the connectivity difference between two groups, with age, gender, ed-




Fig. 6. Validating cluster-level inference at low smoothness by comparing the theoretical cluster-size threshold with permutation-based empirical threshold at FWER 0.05 with different clustering-defining thresholds (from 4.5 to 5.9). The methods are tested in 2 datasets (top: Cambridge; bottom: SWU) under 2 different smoothness levels (0mm and 2mm smoothing). (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

ucation year, head motion (mean frame-wise displacement) being nuisance covariates. The most significant p -value among all functional connectivities was $p = 5.5 \times 10^{-11}$ in this analysis. However, the Bonferroni correction, FDR-BH and FDR-BY approaches cannot detect any significant connectivities (FWER or FDR at 0.05). This is because Bonferroni correction requires the p -value smaller than $p = 4.4 \times 10^{-11}$, and both FDR-BH and FDR-BY approaches require the p -values smaller than the data-driven threshold. See the Manhattan plot for details (Fig. 9).

The p -value threshold of peak-level inference approach was $p = 9.1 \times 10^{-10}$ (connectivity-wise FWER=0.05). A total of 114 altered functional connectivities were found (Fig. 10, left). We also applied cluster-level inference approach to identify significant FC clusters (CDT $p = 3 \times 10^{-7}$ ($Z=5$) and cluster-size FWER=0.05). A total of 12,388 functional connectivities were found with p -values smaller than the applied CDT, and they formed 117 FC clusters. The largest one contains 2247 functional connectivities. Finally, 10 largest FC clusters survived the cluster-size FWER 0.05 threshold (Fig. 10, right).

Almost all the significant FCs in peak-level inference form FC clusters in the cluster-level inference. We could see that, although billions of hypothesis tests were performed and tens of thousands of functional connectivities were found, the results obtained by the cluster-level inference are very structured, thus, easy to be reported (Fig. 10, right). The identified FC clusters can be used in subsequent analysis in several ways. For example, we can calculate the mean functional connectivity within each FC clusters, and use prediction models to classify patients and controls in a new dataset. For patients, we can also test whether the mean functional

connectivity of each FC cluster is associated with the depression symptom severity scores.

3.5. Simulation-based validation for power analysis

Fig. 11 shows the relationship between sample size and power estimated by two methods under two combination of parameters: effect size $\gamma = 0.2, 0.4, 0.6$ and smoothness FWHM=3,4,5,6 voxels. The estimation error (mean squared error) shown on the top of each figure is very low. Therefore, the proposed method can estimate power accurately, and this proposed framework can save a considerable amount of time to generate power curves.

3.6. The power of a future brain-wide association study on MDD

We show an example of how to perform a power analysis to estimate the minimum required sample size for a BWAS. In this example, we will analyze the power of BWAS on MDD using the results of the above study. The aim is to estimate the minimum required sample size to find at least one altered functional connectivities. Base on the above study, the most significant functional connectivities is $p = 5.5 \times 10^{-11}$, corresponding to an effect size of $\gamma = 0.28$. Assuming that in the new dataset, this functional connectivity has a similar effect size, the power under different sample sizes and smoothness levels are estimated and plotted in Fig. 12. Results show that about 80 to 130 subjects are needed to reach 90% power under different smoothness levels.




Fig. 7. Validating peak- and cluster-level inference in *surface-based fMRI data* by comparing the theoretical FWER with permutation-based empirical FWER at 0.05. The methods are tested in 2 datasets (top: Cambridge; bottom: SWU) under 3 different smoothness levels (0, 4, 8 mm smoothing). The estimated FWER is that proportion of permutations in which any significant signals (functional connectivities or FC clusters) are found by the random field theory. Left: results for peak-level inference. Right: results for cluster-level inference with different clustering-defining thresholds (from 4.9 to 5.9). Almost all the results lie in the binomial 95% confidence interval (the dashed line). (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)




Fig. 8. Comparing peak- and cluster-level inference methods with Bonferroni correction and two FDR methods in terms of power (first column) and false discovery rate (second column) across different levels of effect size and smoothness levels using the Cambridge dataset. Left: power curves of different approaches under different smoothness levels. Right: FDR curves of different approaches under different smoothness levels. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)




Fig. 9. Manhattan plot of altered functional connectivities in major depression disorder ($p < 10^{-5}$ only). Each point represents a functional connectivity grouped by the 94 cerebrum regions of the AAL2 template. Bonferroni correction, FDR-BH and FDR-BY fail to identify any significant connections, while both peak- and cluster-level inference approaches identified many altered connectivities. Their brain locations are shown in the next figure. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)




Fig. 10. The altered functional connectivities in major depression disorder identified by peak-level inference (left) and cluster-level inference (right). For peak-level inference, the connectivity-wise FWER is 0.05, which corresponds to uncorrected p -value $= 9 \times 10^{-10}$. For cluster-level inference, the CDT is $Z=5$ ($p=3 \times 10^7$) and cluster-size FWER threshold is 0.05. Abbreviations of regions are listed in Supplement Table G.2. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

4. Discussion

Our proposed method can accurately control both connectivity-wise FWER and connectivity-extent FWER, as demonstrated by comparing with the empirical FWER obtained from two real datasets. To the best of our knowledge, BWAS is the first method to use the random field theory to analyze the voxel-wise functional connectome. Random field theory makes some assumptions of data. Eklund et al. (2016) recently reported that random field theory could lead to inflated false-positive rate in task-activation analysis, particularly when the CDT is low ($p = 0.01$). This failure is well known since the choice of low CDT violates the assumptions of the original theory (Friston et al., 1994). However, when the CDT is selected appropriately (e.g. $p = 0.001$), FWER is closer to nominal level (Eklund et al., 2016; Flandin and Friston, 2016; Greve and Fischl, 2017). Another article (Flandin and Friston, 2016) has also pointed out that the random field theory can provide accept-

able FWER control when using two-sample t -test instead of one-sample t -test and resampling the data closer to the original image resolution. In our analysis, we have demonstrated that the random field theory is valid for both volume- and surface-based resting-state fMRI data under different smoothness. Particularly, the CDT in cluster-level inference should be high enough ($|Z| > 5$ for moderate or large smoothness and $|Z| > 4.5$ for low smoothness).

Importantly, our method is computationally efficient. It is a fully parametric approach which is not based on any simulation or permutation. Although non-parametric approaches can also perform multiple correction and power analysis (Nichols and Holmes, 2002), they are extremely slow in voxel-wise analysis by the necessity of calculating billions of statistical tests many times. Our empirical studies show that our approach is usually at least N times faster than the non-parametric permutation approaches, where N is the number of permutations performed. The reason is that, although the subject-level brain network can be




Fig. 11. Comparing the theoretical power analysis method (red line) with the simulation result (blue line). Each figure shows the relationship between estimated power and sample size. From the left to the right, the effect sizes are 0.2, 0.4 and 0.6. From the top to the bottom, the FWHMs are 3 to 6 voxels. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 12. Power of detecting at least one altered functional connectivity in major depression disorder under different sample sizes and smoothness levels estimated by the proposed approach. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

computed only once in non-parametric permutations, the fitting of connexel-wise GLM is usually much slower than network construction, thus, it dominates the computation time. In addition, parallelization of permutations will not save much time, because the transmission speed of large data between processors is very slow.

Many possible extensions and improvements of the current framework can be developed in the future. First, the functional connectivities identified by massive univariate statistical tests approach may not be predictive, e.g., in a case-control study, the identified connectivities may not be able to classify patients and controls. A directly construction of connexel-wise prediction model is also not practical, since the model constructed on a few hundred subjects and billion of features usually has a large variance. Meanwhile, the optimization of model parameters become very difficult in this ultra-high dimensional feature space. One possible way to solve this problem is to adopt the current BWAS frame-

work into sure independence screening (SIS) approach (Fan and Lv, 2008; Fan et al., 2009; 2010). Second, this framework can be extended to task fMRI analysis to identify network configuration changes (e.g. Lohmann et al., 2016). It can support either single subject analysis or group analysis provided that the task experiment is in block design and the length of each trial is long enough to enable network construction. Third, the cluster-level inference proposed here controls the FWER of cluster size. An alternative method of controlling the FDR of cluster size was proposed in task-activation studies (Chumbley and Friston, 2009; Chumbley et al., 2010), which can be easily adopted here. Forth, the estimation of subject-level functional network is based on the Pearson correlation between pairwise BOLD signal time series in the current framework, which may be suboptimal (Westfall and Yarkoni, 2016; Bellec et al., 2008; Sahib et al., 2016). Therefore, a better approach for constructing a functional network at the voxel level should be designed and validated in the future (Narayan and Allen, 2016; Bickel and Levina, 2008). Fifth, with the higher volume of available data, statistical methods for combining BWAS results from multiple imaging centers are needed. In BWAS, integrating results from different datasets has been shown to greatly reduce the false-positive rate and increase sensitivity (Cheng et al., 2015a; 2015b; 2016). However, the sample heterogeneity introduced by different sources, such as different data acquisition pipelines, population stratification, and genetic background, may make the traditional meta-analysis methods used in our previous studies suboptimal.

In this paper, we develop a rigorous statistical framework for BWAS. Both peak- and cluster-level inferences are introduced for the analysis of voxel-wise functional connectomes, and the random field theory is developed to control FWER and estimate statistical power. We believe that this method will be very useful for the neuroimaging fields in the context of understanding the brain connectome.

Acknowledgments

Jianfeng Feng is supported by the National High Technology Research and Development Program of China (No. 2015AA020507),

the Key Program of [National Natural Science Foundation of China](#) (No. [91230201](#)), International (Regional) Collaborative and Exchange Program of National Natural Science (No. 71661167002), the Key Project of Shanghai Science and Technology Innovation Plan (No. 15JC1400101), and the Shanghai Soft Science Research Program (No. 15692106604), and the National Centre for Mathematics and Interdisciplinary Sciences (NCMIS) of the Chinese Academy of Sciences. Jianfeng Feng is a Royal Society Wolfson Research Merit Award holder. Lin Wan is supported by the NSFC grants (Nos. [11571349](#) and [11201460](#)), the NCMIS of the CAS, the Youth Innovation Promotion Association of the CAS, and the Strategic Priority Research Program of the CAS (XDB13040600). Wenlian Lu is jointly supported by the Natural Sciences Foundation of China under Grant No. [61673119](#), the Shanghai Science and Technology Innovation Project under grant No. 17411953500, the Laboratory of Mathematics for Nonlinear Science, Fudan University, and the Shanghai Key Laboratory for Contemporary Applied Mathematics, Fudan University. Wenlian Lu would thank Prof. Ling Yang from School of Mathematical Sciences, Fudan University, for valuable discussions and suggestions. Wei Cheng is supported by grants from the National Natural Sciences Foundation of China (No. [81701773](#)), Sponsored by Shanghai Sailing Program (No. 17YF1426200) and the Research Fund for the Doctoral Program of Higher Education of China (No. 2017M610226).

Appendix A. Image acquisition and preprocessing

Only publicly available data are used in this article. Resting-state fMRI data are collected from two imaging sites: (1) 197 samples from the Cambridge dataset in 1000 Functional Connectomes Project (1000 FCP) ([Biswal et al., 2010](#)) (http://fcon_1000.projects.nitrc.org/fcpClassic/FcpTable.html); (2) 222 subjects from the Southwest University dataset in International Data-sharing Initiative (IDNI) (http://fcon_1000.projects.nitrc.org/indirect/southwestuni_qiu_index.html). All subjects are normal people. As Southwest University dataset is a longitudinal dataset, only subjects who scanned at the first time are used in this paper.

All the data collected are subject to their local ethics review boards, the experiments and the dissemination of the anonymized data are approved. The detailed data acquisition methods may be found in the respective websites and papers. The data were preprocessed using SPM12 ([Penny et al., 2011](#)) and Data Processing and Analysis for Brain Imaging (DPABI) ([Yan et al., 2016](#)). For each individual, the preprocessing steps included discarding the first 10 time points, slice timing correction, motion correction, coregistering the functional image to individual T1 structure image, segmenting structure images and DARTEL registration ([Ashburner, 2007](#)), regressing out nuisance covariates including 24 head motion parameters ([Friston et al., 1996](#)), white matter signals, cerebrospinal fluid signals, temporal filtering (0.01–0.1 Hz), normalizing to standard space of voxel size $3 \times 3 \times 3 \text{ mm}^3$ by DARTEL, and smoothing by a 3D Gaussian kernel with FWHM = 0, 2, 4, 6, 8, 10, 12 mm. Finally, all the images are manually checked by experts to ensure preprocessing quality. Images that are not successfully preprocessed are discarded in our analysis.

The surface-based fMRI data are preprocessed using Connectome workbench. For each volume-based fMRI data in the Cambridge and Southwest University dataset, we map it to the Conte69 surface-based atlas (http://brainvis.wustl.edu/wiki/index.php/Caret:Atlases/Conte69_Atlas) using the command 'wb_command -volume-to-surface-mapping'. Each fMRI images are then smoothed by a 2D Gaussian kernel with FWHM=0, 4, 8 mm using the command 'wb_command cifti-smoothing'. Finally, the smoothness of each image is estimated by the command 'wb_command -cifti-estimate-fwhm'. The surface area of Conte69

is estimated using the command 'wb_command -surface-vertex-areas', which is used in the random field theory.

Appendix B. Calculating the intrinsic volume and Gaussian EC-density

To perform peak-level and cluster-level inference, we should calculate the 0- to 3-dimensional intrinsic volume and the 0- to 6-dimensional EC-densities for the Gaussian random field.

Let P be the number of voxels, E_x (or E_y, E_z) be number of x (or y, z)-direction edges (two adjacent voxels), F_{xy} (or F_{yz}, F_{xz}) be number of xy (or yz, xz)-direction surface (four adjacent voxels), and C be the number of cubes (eight adjacent voxels). The r_x (or r_y, r_z) be the resel size of x (or y, z)-direction, which is defined as the voxel size divided by FWHM (in mm). The 0–3 dimensional intrinsic volume of S can be calculated as:

$$\begin{aligned} u_0(S) &= P - (E_x + E_y + E_z) + (F_{yz} + F_{xz} + F_{xy}) - C \\ u_1(S) &= (E_x - F_{xy} - F_{xz} + C)r_x + (E_y - F_{xy} - F_{yz} + C)r_y \\ &\quad + (E_z - F_{xz} - F_{yz} + C)r_z \\ u_2(S) &= (F_{xy} - C)r_x r_y + (F_{xz} - C)r_x r_z + (F_{yz} - C)r_y r_z \\ u_3(S) &= Cr_x r_y r_z \end{aligned}$$

The above calculation has been implement in SPM package as *spm_resels_vol* function. Two other methods also work well in practice. One is to replace the original space with an equal volume ball, as implement in the *fmrstat* package, the other is to use a linear regression model ([Adler et al., 2017](#)), which do not need the knowledge of spatial smoothness. In whole-brain BWAS, for peak-level inference, the $u_i(P)$ and $u_i(Q)$ are the same. As there are $p(p-1)/2$ functional connectivities across p voxels, we divided the estimated intrinsic volume by $\sqrt{2}$, thus, the highest order term, $u_3(P) \times u_3(Q)$, will approximate the total number of functional connectivities (in resel) in the brain.

The 0- to 6-dimensional EC-densities for Gaussian random field at t are:

$$\begin{aligned} \rho_0(t) &= 1 - \Phi(t) \\ \rho_1(t) &= (4 \ln 2)^{\frac{1}{2}} (2\pi)^{-1} e^{-\frac{t^2}{2}} \\ \rho_2(t) &= (4 \ln 2) (2\pi)^{-\frac{3}{2}} t e^{-\frac{t^2}{2}} \\ \rho_3(t) &= (4 \ln 2)^{\frac{3}{2}} (2\pi)^{-2} (t^2 - 1) e^{-\frac{t^2}{2}} \\ \rho_4(t) &= (4 \ln 2)^2 (2\pi)^{-\frac{5}{2}} (t^3 - 3t) e^{-\frac{t^2}{2}} \\ \rho_5(t) &= (4 \ln 2)^{\frac{5}{2}} (2\pi)^{-3} (t^4 - 6t^2 + 3) e^{-\frac{t^2}{2}} \\ \rho_6(t) &= (4 \ln 2)^3 (2\pi)^{-\frac{7}{2}} (t^5 - 10t^3 + 15t) e^{-\frac{t^2}{2}} \end{aligned}$$

where $\Phi(\bullet)$ is the cumulative distribution function of standard normal distribution.

Appendix C. Estimating the smoothness of the fMRI images

The true smoothness of fMRI images is usually larger than the applied smoothness. Therefore, an accurate estimation of smoothness is critical for Gaussian random field theory. The following approach is used to estimate the smoothness of 3D or 2D images ([Hagler et al., 2006](#)):

$$\text{FWHM} = dv \sqrt{\frac{-2 \ln 2}{\ln(1 - \frac{\text{var}(ds)}{2\text{var}(s)})}}$$

where dv is the average inter-neighbor distance of voxels or vertices, $\text{var}(ds)$ is the variance of inter-neighbors differences, and $\text{var}(s)$ is the overall variance of the values at each voxels or vertices. The FWHM of fMRI image is the average smoothness of the 3D or 2D images across all time points.

Appendix D. Proof of Eq. (1) in the main text

Using the property of d -dimensional intrinsic volume (Taylor and Worsley, 2008)

$$u_d(\mathcal{P} \times \mathcal{Q}) = \sum_{k=0}^d u_k(\mathcal{P}) u_{d-k}(\mathcal{Q})$$

When $d > P$, $u_d(\mathcal{P}) = 0$ and $d > Q$, $u_d(\mathcal{Q}) = 0$. It is easy to conclude that

$$\begin{aligned} \sum_{d=0}^{P+Q} \mu_d(\mathcal{P} \times \mathcal{Q}) &= \sum_{d=0}^{P+Q} \sum_{k=0}^d u_k(\mathcal{P}) u_{d-k}(\mathcal{Q}) \\ &= \sum_{i=0}^P \sum_{j=0}^Q \mu_i(\mathcal{P}) \mu_j(\mathcal{Q}) \end{aligned}$$

Appendix E. Proof of Eq. (2) in the main text

We begin by considering the normal transformation of t -statistic of BWAS map, which makes the $\hat{\sigma}_{ij}^2$ fixed as $\hat{\sigma}_0^2$ (Worsley et al., 1992; 1996). Therefore, the test statistic becomes

$$\begin{aligned} Z_{ij} &= \frac{\mathbf{c}\hat{\mathbf{b}}_{ij}}{\sqrt{\mathbf{c}(X'X)^{-1}\mathbf{c}'\hat{\sigma}_0^2}} \\ &= \frac{\mathbf{c}(X'X)^{-1}X'Y_{ij}}{\sqrt{\mathbf{c}(X'X)^{-1}\mathbf{c}'\hat{\sigma}_0^2}} \\ &= \sum_{s=1}^n w^{(s)} z_{ij}^{(s)} \end{aligned}$$

where $w^{(s)}$ is the s -th element of row vector $\frac{\mathbf{c}(X'X)^{-1}X'Y_{ij}}{\sqrt{\mathbf{c}(X'X)^{-1}\mathbf{c}'\hat{\sigma}_0^2}}$, which only depends on the subjects, and $z_{ij}^{(s)}$ is Fisher's Z transformed correlation coefficient.

Let $M^{(s)}(p) = (M_1^{(s)}(p), \dots, M_{\nu^{(s)}}^{(s)}(p))'$, $p \in \mathcal{P} \subset \mathbb{R}^P$ and $N^{(s)}(q) = (N_1^{(s)}(q), \dots, N_{\nu^{(s)}}^{(s)}(q))$, $q \in \mathcal{Q} \subset \mathbb{R}^Q$ be two vectors of $\nu^{(s)}$ independent and stationary Gaussian random fields with mean zeros and variance one. The index s denotes subjects, and the $\nu^{(s)}$ can be treated as the number of time points, while p, q are the coordinates of three-dimensional Euclidean space. The $(P+Q)$ -dimensional cross-correlation random field $R^{(s)}(p, q)$ is defined as follows (Cao et al., 1999):

$$R^{(s)}(p, q) = \frac{M^{(s)}(p)'N^{(s)}(q)}{\sqrt{M^{(s)}(p)'M^{(s)}(p)N^{(s)}(q)'N^{(s)}(q)}}$$

In BWAS, the cross-correlation field is generated by calculating Pearson correlation coefficients between pairwise voxel time series. Next, the element-wise Fisher's Z transformation transforms this cross-correlation random field to a six-dimensional 'Gaussianized' random field as:

$$Z^{(s)}(p, q) = \frac{1}{2} \log \left[\frac{1 + R^{(s)}(p, q)}{1 - R^{(s)}(p, q)} \right]$$

It has mean zero and variance $\frac{1}{\nu^{(s)}-3}$ (Kenney, 2013). Our test statistic $Z_{ij}(p, q)$ forms a weighted sum of Fisher's Z transformed cross-correlation random field $Z(p, q)$ as:

$$Z(p, q) = \sum_{s=1}^n w^{(s)} Z^{(s)}(p, q)$$

The random field $Z(p, q)$ is a 'Gaussianized' random field with mean zero and variance one. Therefore, we can use Eq. (1) in the

main text to approximate its maximum distribution at high threshold:

$$\begin{aligned} P(\max Z(p, q) > z_0) &\approx \mathbb{E}(EC) = \sum_{d=0}^{P+Q} \mu_d(\mathcal{P} \times \mathcal{Q}) \rho_d^Z(z_0) \\ &= \sum_{i=0}^P \sum_{j=0}^Q \mu_i(\mathcal{P}) \mu_j(\mathcal{Q}) \rho_{i+j}^Z(z_0) \end{aligned} \quad (E.1)$$

The EC-densities for the Gaussian random field $\rho_d^Z(z_0)$ in any dimensions can be expressed as (Adler and Taylor, 2009):

$$\rho_d^Z(z_0) = (2\pi)^{-\frac{d+1}{2}} |\Lambda|^{\frac{d}{2(P+Q)}} e^{-\frac{z_0^2}{2}} \sum_{j=0}^{\lfloor \frac{d-1}{2} \rfloor} (-1)^j \frac{(2j)!}{j!2^j} \binom{d-1}{2j} z_0^{d-1-2j}$$

where D is the highest dimension of $Z(p, q)$. The $|\Lambda| = |\text{Var}(\dot{Z}(p, q))|$ is the determinant of the variance-covariance matrix of the partial derivative of $Z(p, q)$. The $|\Lambda|$ can be replaced by FWHM_Z , the Full Width at Half Maximum (FWHM) of the random field Z averaged across all dimensions, using the equation:

$$\text{FWHM}_Z = (4 \log 2)^{\frac{1}{2}} |\Lambda|^{-\frac{1}{2D}}$$

and FWHM_Z is a corrected smoothness parameter, which can be calculated as:

$$\begin{aligned} \text{FWHM}_Z &= \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)}-3} \text{FWHM}_{M^{(s)}}^{-2} \right)^{-\frac{P}{2(P+Q)}} \\ &\quad \times \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)}-3} \text{FWHM}_{N^{(s)}}^{-2} \right)^{-\frac{Q}{2(P+Q)}} \end{aligned}$$

where $\text{FWHM}_{M^{(s)}}$ and $\text{FWHM}_{N^{(s)}}$ are the average FWHM of the random field vectors $M^{(s)}(p)$ and $N^{(s)}(q)$ across three dimensions. The proof of this equation is given in the next section. Finally, Eq. (2) in the main text is used in the peak-level inference:

$$\begin{aligned} P(\max Z(p, q) > z_0) &\approx \sum_{i=0}^P \sum_{j=0}^Q \mu_i(\mathcal{P}) \mu_j(\mathcal{Q}) \frac{(2\pi)^{-\frac{i+j+1}{2}} (4 \log 2)^{\frac{i+j}{2}}}{\text{FWHM}_Z^{i+j}} \\ &\quad \times e^{-\frac{z_0^2}{2}} \sum_{k=0}^{\lfloor \frac{i+j-1}{2} \rfloor} (-1)^k \frac{(2k)!}{k!2^k} \binom{i+j-1}{2k} z_0^{d-1-2k} \end{aligned} \quad (E.2)$$

We can see that FWHM_Z is a function of the number of time points $\nu^{(s)}$, the $\text{FWHM}_{M^{(s)}}$ and $\text{FWHM}_{N^{(s)}}$ of the individual fMRI data. Assuming that the length of scanning time and image smoothness is the same for every subject in a study. Denoting them as ν and FWHM , the formula for calculating FWHM_Z reduces to:

$$\begin{aligned} \text{FWHM}_Z &= \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu-3} \text{FWHM}^{-2} \right)^{-\frac{P}{2(P+Q)}} \\ &\quad \times \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu-3} \text{FWHM}^{-2} \right)^{-\frac{Q}{2(P+Q)}} \\ &= \text{FWHM} \sqrt{\nu-3} \left[\left(\frac{\mathbf{c}(X'X)^{-1}X'}{\sqrt{\mathbf{c}(X'X)\mathbf{c}'\hat{\sigma}_0^2}} \right) \left(\frac{\mathbf{c}(X'X)^{-1}X'}{\sqrt{\mathbf{c}(X'X)\mathbf{c}'\hat{\sigma}_0^2}} \right)' \right]^{-\frac{1}{2}} \\ &= \text{FWHM} \sqrt{(\nu-3)\sigma_0^2} \\ &= \text{FWHM} \end{aligned}$$

where we treat the sample variance σ_0^2 as the theoretical variance $\frac{1}{\nu-3}$. This suggests that the smoothness of random field $Z(p, q)$ equals the original image smoothness. A series of non-linear transformations will maintain the original smoothness of images. The scanning time does not influence this equation.

In practice, for volume-based BWAS, we found that Eq. (1) usually provide a slightly conservative estimation of FWER-corrected threshold. Therefore, for volume-based BWAS, we modify it as:

$$\alpha = P(\max Z(p, q) > z_0) \approx \mathbb{E}(EC) \approx \mu_2(\mathcal{P})\mu_2(\mathcal{Q})\rho_6^Z(z_0) \quad (E.3)$$

For surface-based BWAS, we use:

$$\alpha = P(\max Z(p, q) > z_0) \approx \mathbb{E}(EC) = \sum_{i=0}^2 \sum_{j=0}^2 \mu_i(\mathcal{P})\mu_j(\mathcal{Q})\rho_{i+j}^Z(z_0) \quad (E.4)$$

If we only analyse the functional connectivities between subcortical voxels \mathcal{P} and cortical vertices \mathcal{Q} , we use:

$$\alpha = P(\max Z(p, q) > z_0) \approx \mathbb{E}(EC) = \sum_{i=0}^3 \sum_{j=0}^2 \mu_i(\mathcal{P})\mu_j(\mathcal{Q})\rho_{i+j}^Z(z_0) \quad (E.5)$$

Proof of the equation for calculating FWHM_Z

Let $Var(M^{(s)}(p)) = \Lambda_{M_s}$ and $Var(N^{(s)}(q)) = \Lambda_{N_s}$, then according to Lemma 4.2 in Cao et al. (1999),

$$\frac{\partial R^{(s)}(p, q)}{\partial p} \stackrel{D}{=} (1 - R^{(s)}(p, q)^2)^{\frac{1}{2}} a_s^{-\frac{1}{2}} (\Lambda_{M_s})^{\frac{1}{2}} z_M^{(s)}$$

and

$$\frac{\partial R^{(s)}(p, q)}{\partial q} \stackrel{D}{=} (1 - R^{(s)}(p, q)^2)^{\frac{1}{2}} a_s^{-\frac{1}{2}} (\Lambda_{N_s})^{\frac{1}{2}} z_N^{(s)}$$

where $a_s \sim \chi_{\nu^{(s)}}^2$, $z_M^{(s)} \sim N(0, I_{p,p})$, $z_N^{(s)} \sim N(0, I_{q,q})$ and independent of $R^{(s)}(p, q)$, and $\stackrel{D}{=}$ means equal in distribution. Then, after Fisher's Z transformation, we have

$$\frac{\partial Z^{(s)}(p, q)}{\partial p} \stackrel{D}{=} (1 - R^{(s)}(p, q)^2)^{-\frac{1}{2}} a_s^{-\frac{1}{2}} (\Lambda_{M_s})^{\frac{1}{2}} z_M^{(s)}$$

and

$$\frac{\partial Z^{(s)}(p, q)}{\partial q} \stackrel{D}{=} (1 - R^{(s)}(p, q)^2)^{-\frac{1}{2}} a_s^{-\frac{1}{2}} (\Lambda_{N_s})^{\frac{1}{2}} z_N^{(s)}$$

Then,

$$\frac{\partial Z(p, q)}{\partial p} \stackrel{D}{=} \sum_{s=1}^n w^{(s)} (1 - R^{(s)}(p, q)^2)^{-\frac{1}{2}} a_s^{-\frac{1}{2}} (\Lambda_{M_s})^{\frac{1}{2}} z_M^i$$

and

$$\frac{\partial Z(p, q)}{\partial q} \stackrel{D}{=} \sum_{s=1}^n w^{(s)} (1 - R^{(s)}(p, q)^2)^{-\frac{1}{2}} a_s^{-\frac{1}{2}} (\Lambda_{N_s})^{\frac{1}{2}} z_N^i$$

Since

$$Var(\dot{Z}(p, q)) = \begin{pmatrix} \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial p} \frac{\partial Z^{(s)'}(p, q)}{\partial p} \right] & \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial p} \frac{\partial Z^{(s)'}(p, q)}{\partial q} \right] \\ \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial q} \frac{\partial Z^{(s)'}(p, q)}{\partial p} \right] & \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial q} \frac{\partial Z^{(s)'}(p, q)}{\partial q} \right] \end{pmatrix}$$

and

$$\mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial p} \frac{\partial Z^{(s)'}(p, q)}{\partial q} \right] = \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial q} \frac{\partial Z^{(s)'}(p, q)}{\partial p} \right] = 0$$

and

$$\begin{aligned} & \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial p} \frac{\partial Z^{(s)'}(p, q)}{\partial p} \right] \\ &= \sum_{s=1}^n (w^{(s)})^2 \mathbb{E}[(1 - R^{(s)}(p, q)^2)^{-1}] \mathbb{E}[a_s^{-1}] \Lambda_{M_s} \\ & \mathbb{E} \left[\frac{\partial Z^{(s)}(p, q)}{\partial q} \frac{\partial Z^{(s)'}(p, q)}{\partial q} \right] \\ &= \sum_{s=1}^n (w^{(s)})^2 \mathbb{E}[(1 - R^{(s)}(p, q)^2)^{-1}] \mathbb{E}[a_s^{-1}] \Lambda_{N_s} \end{aligned}$$

The expectations in the above equations are

$$\mathbb{E}[a_s^{-1}] = \frac{1}{\nu^{(s)} - 2}$$

$$\mathbb{E}[(1 - R^{(s)}(p, q)^2)^{-1}] = \frac{\nu^{(s)} - 2}{\nu^{(s)} - 3}$$

Finally we get

$$Var(\dot{Z}(p, q)) = \begin{pmatrix} \sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)} - 3} \Lambda_{M_s} & 0 \\ 0 & \sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)} - 3} \Lambda_{N_s} \end{pmatrix}_{6 \times 6}$$

Substituting the variance covariance matrix of partial derivative of the random field by the FWHM using its relationship with $|\Lambda|$, we could get:

$$\begin{aligned} & \left(\frac{FWHM_Z}{(4 \ln 2)^{\frac{1}{2}}} \right)^{-2(P+Q)} \\ &= \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)} - 3} \frac{FWHM_{M^{(s)}}^{-2}}{(4 \ln 2)^{\frac{1}{2}}} \right)^P \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)} - 3} \frac{FWHM_{N^{(s)}}^{-2}}{(4 \ln 2)^{\frac{1}{2}}} \right)^Q \end{aligned}$$

thus

$$\begin{aligned} FWHM_Z &= \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)} - 3} FWHM_{M^{(s)}}^{-2} \right)^{-\frac{P}{2(P+Q)}} \\ & \times \left(\sum_{s=1}^n \frac{(w^{(s)})^2}{\nu^{(s)} - 3} FWHM_{N^{(s)}}^{-2} \right)^{-\frac{Q}{2(P+Q)}} \end{aligned}$$

Supplementary material

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.media.2018.03.014.

References

Adler, R.J., 1981. *The Geometry of Random Fields*, Vol. 62. SIAM.
 Adler, R.J., Taylor, J.E., 2009. *Random Fields and Geometry*. Springer Science & Business Media.
 Adler, R.J., Bartz, K., Kou, S.C., Monod, A., 2017. Estimating thresholding levels for random fields via Euler characteristics. arXiv preprint arXiv:1704.08562
 Ashburner, J., 2007. A fast diffeomorphic image registration algorithm. *NeuroImage* 38 (1), 95–113.
 Ashburner, J., Friston, K.J., 2000. Voxel-based morphometry—the methods. *NeuroImage* 11 (6), 805–821.
 Beckmann, C.F., Smith, S.M., 2004. Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans. Med. Imaging* 23 (2), 137–152.
 Bellec, P., Marrelec, G., Benali, H., 2008. A bootstrap test to investigate changes in brain connectivity for functional MRI. *Stat. Sin.* 1253–1268.
 Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B (Methodol.)* 289–300.

- Benjamini, Y., Yekutieli, D., 2001. The control of the false discovery rate in multiple testing under dependency. *Ann. Stat.* 1165–1188.
- Bickel, P.J., Levina, E., 2008. Regularized estimation of large covariance matrices. *Ann. Stat.* 199–227.
- Biswal, B.B., Mennes, M., Zuo, X.-N., Gohel, S., Kelly, C., Smith, S.M., Beckmann, C.F., Adelstein, J.S., Buckner, R.L., Colcombe, S., et al., 2010. Toward discovery science of human brain function. *Proc. Natl. Acad. Sci.* 107 (10), 4734–4739.
- Cao, J., 1999. The size of the connected components of excursion sets of χ^2 , t and F fields. *Adv. Appl. Probab.* 579–595.
- Cao, J., Worsley, K., et al., 1999. The geometry of correlation fields with an application to functional connectivity of the brain. *Ann. Appl. Probab.* 9 (4), 1021–1057.
- Cheng, W., Palaniyappan, L., Li, M., Kendrick, K.M., Zhang, J., Luo, Q., Liu, Z., Yu, R., Deng, W., Wang, Q., et al., 2015a. Voxel-based, brain-wide association study of aberrant functional connectivity in schizophrenia implicates thalamocortical circuitry. *NPJ Schizophr.* 1.
- Cheng, W., Rolls, E.T., Gu, H., Zhang, J., Feng, J., 2015b. Autism: reduced connectivity between cortical areas involved in face expression, theory of mind, and the sense of self. *Brain* 138 (5), 1382–1393. aww051
- Cheng, W., Rolls, E.T., Qiu, J., Liu, W., Tang, Y., Huang, C.-C., Wang, X., Zhang, J., Lin, W., Zheng, L., et al., 2016. Medial reward and lateral non-reward orbitofrontal cortex circuits change in opposite directions in depression. *Brain* 139 (12), 3296–3309. aww255
- Chumbley, J., Worsley, K., Flandin, G., Friston, K., 2010. Topological FDR for neuroimaging. *NeuroImage* 49 (4), 3057–3064.
- Chumbley, J.R., Friston, K.J., 2009. False discovery rate revisited: FDR and topological inference using gaussian random fields. *NeuroImage* 44 (1), 62–70.
- Desmond, J.E., Glover, G.H., 2002. Estimating sample size in functional MRI (fMRI) neuroimaging studies: statistical power analyses. *J. Neurosci. Methods* 118 (2), 115–128.
- Eklund, A., Andersson, M., Josephson, C., Johannesson, M., Knutsson, H., 2012. Does parametric fMRI analysis with SPM yield valid results?—An empirical study of 1484 rest datasets. *NeuroImage* 61 (3), 565–578.
- Eklund, A., Nichols, T.E., Knutsson, H., 2016. Cluster failure: why fMRI inferences for spatial extent have inflated false-positive rates. *Proc. Natl. Acad. Sci.* 113 (28), 7900–7905. 201602413
- Fan, J., Lv, J., 2008. Sure independence screening for ultrahigh dimensional feature space. *J. R. Stat. Soc. Ser. B (Stat. Methodol.)* 70 (5), 849–911.
- Fan, J., Samworth, R., Wu, Y., 2009. Ultrahigh dimensional feature selection: beyond the linear model. *J. Mach. Learn. Res.* 10, 2013–2038. Sep
- Fan, J., Song, R., et al., 2010. Sure independence screening in generalized linear models with NP-dimensionality. *Ann. Stat.* 38 (6), 3567–3604.
- Flandin, G., Friston, K.J., 2016. Analysis of family-wise error rates in statistical parametric mapping using random field theory. *Hum. Brain Mapp.*
- Friston, K.J., Williams, S., Howard, R., Frackowiak, R.S., Turner, R., 1996. Movement-related effects in fMRI time-series. *Magn. Reson. Med.* 35 (3), 346–355.
- Friston, K.J., Worsley, K.J., Frackowiak, R., Mazziotta, J.C., Evans, A.C., 1994. Assessing the significance of focal activations using their spatial extent. *Hum. Brain Mapp.* 1 (3), 210–220.
- Glasser, M.F., Smith, S.M., Marcus, D.S., Andersson, J.L., Auerbach, E.J., Behrens, T.E., Coalson, T.S., Harms, M.P., Jenkinson, M., Moeller, S., et al., 2016. The human connectome project's neuroimaging approach. *Nat. Neurosci.* 19 (9), 1175–1187.
- Greve, D.N., Fischl, B., 2017. False positive rates in surface-based anatomical analysis. *NeuroImage*.
- Hagler, D.J., Saygin, A.P., Sereno, M.I., 2006. Smoothing and cluster thresholding for cortical surface-based group analysis of fMRI data. *NeuroImage* 33 (4), 1093–1103.
- Hayasaka, S., Nichols, T.E., 2003. Validating cluster size inference: random field and permutation methods. *NeuroImage* 20 (4), 2343–2356.
- Hayasaka, S., Peiffer, A.M., Hugenschmidt, C.E., Laurienti, P.J., 2007. Power and sample size calculation for neuroimaging studies by non-central random field theory. *NeuroImage* 37 (3), 721–730.
- Joyce, K.E., Hayasaka, S., 2012. Development of powermap: a software package for statistical power calculation in neuroimaging studies. *Neuroinformatics* 10 (4), 351–365.
- Kenney, J.F., 2013. *Mathematics of Statistics*. D. Van Nostrand Company Inc, Toronto; Princeton; New Jersey; London; New York Affiliated East-West Press Pvt-Ltd; New Delhi.
- Kim, J., Wozniak, J.R., Mueller, B.A., Shen, X., Pan, W., 2014. Comparison of statistical tests for group differences in brain functional networks. *NeuroImage* 101, 681–694.
- Lohmann, G., Stelzer, J., Zuber, V., Buschmann, T., Margulies, D., Bartels, A., Scheffler, K., 2016. Task-related edge density (TED) – a new method for revealing dynamic network formation in fMRI data of the human brain. *PLoS One* 11 (6), e0158185.
- Miller, K.L., Alfaro-Almagro, F., Bangerter, N.K., Thomas, D.L., Yacoub, E., Xu, J., Bartsch, A.J., Jbabdi, S., Sotiropoulos, S.N., Andersson, J.L., et al., 2016. Multimodal population brain imaging in the UK biobank prospective epidemiological study. *Nat. Neurosci.* 19 (11), 1523.
- Mumford, J.A., Nichols, T.E., 2008. Power calculation for group fMRI studies accounting for arbitrary design and temporal autocorrelation. *NeuroImage* 39 (1), 261–268.
- Narayan, M., Allen, G.I., 2016. Mixed effects models for resampled network statistics improves statistical power to find differences in multi-subject functional connectivity. *Front. Neurosci.* 10, 15016.
- Nichols, T.E., Holmes, A.P., 2002. Nonparametric permutation tests for functional neuroimaging: a primer with examples. *Hum. Brain Mapp.* 15 (1), 1–25.
- Nosko, V., 1969. Local structure of gaussian random fields in the neighborhood of high-level shines. *Dokl. Akad. Nauk SSSR* 189, 714–717.
- Penny, W.D., Friston, K.J., Ashburner, J.T., Kiebel, S.J., Nichols, T.E., 2011. *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. Academic Press.
- Rolls, E.T., Joliot, M., Tzourio-Mazoyer, N., 2015. Implementation of a new parcellation of the orbitofrontal cortex in the automated anatomical labeling atlas. *NeuroImage* 122, 1–5.
- Sahib, A.K., Mathiak, K., Erb, M., Elshahabi, A., Klamer, S., Scheffler, K., Focke, N.K., Ethofer, T., 2016. Effect of temporal resolution and serial autocorrelations in event-related functional MRI. *Magn. Reson. Med.* 76 (6), 1805–1813.
- Taylor, J., Worsley, K., 2008. Random fields of multivariate test statistics, with applications to shape analysis. *Ann. Stat.* 1–27.
- Westfall, J., Yarkoni, T., 2016. Statistically controlling for confounding constructs is harder than you think. *PLoS One* 11 (3), e0152719.
- Worsley, K.J., 1994. Local maxima and the expected Euler characteristic of excursion sets of χ^2 , f and T fields. *Adv. Appl. Probab.* 13–42.
- Worsley, K.J., Evans, A.C., Marrett, S., Neelin, P., et al., 1992. A three-dimensional statistical analysis for CBF activation studies in human brain. *J. Cereb. Blood Flow Metab.* 12, 900–918.
- Worsley, K.J., Marrett, S., Neelin, P., Vandal, A.C., Friston, K.J., Evans, A.C., et al., 1996. A unified statistical approach for determining significant signals in images of cerebral activation. *Hum. Brain Mapp.* 4 (1), 58–73.
- Yan, C.-G., Wang, X.-D., Zuo, X.-N., Zang, Y.-F., 2016. DPABI: data processing & analysis for (resting-state) brain imaging. *Neuroinformatics* 1–13.
- Yang, J., Zaitlen, N.A., Goddard, M.E., Visscher, P.M., Price, A.L., 2014. Advantages and pitfalls in the application of mixed-model association methods. *Nat. Genet.* 46 (2), 100–106.
- Zalesky, A., Cocchi, L., Fornito, A., Murray, M.M., Bullmore, E., 2012. Connectivity differences in brain networks. *NeuroImage* 60 (2), 1055–1062.
- Zhou, X., Stephens, M., 2012. Genome-wide efficient mixed-model analysis for association studies. *Nat. Genet.* 44 (7), 821–824.